



A new role for pond management in farmland bird conservation



S.R. Davies^{a,*}, C.D. Sayer^a, H. Greaves^a, G.M. Siriwardena^b, J.C. Axmacher^a

^a Pond Restoration Research Group, Environmental Change Research Centre, Department of Geography, University College London, London, UK

^b British Trust for Ornithology, The Nunnery, Thetford, Norfolk IP24 2PU, UK

ARTICLE INFO

Article history:

Received 30 June 2016

Received in revised form 7 September 2016

Accepted 9 September 2016

Available online xxx

Keywords:

Agri-environment schemes

Agro-ecosystems

Avian diversity

Biodiversity decline

Farmland ponds

Habitat heterogeneity

ABSTRACT

Biodiversity declines in agricultural landscapes represent a major conservation challenge. In the UK, some agricultural landscapes contain high pond densities, but many farmland ponds have become terrestrialised since the 1960s, with input of organic material resulting in a decrease in the size and depth of ponds that eventually transform into wet woodland habitats. Pond management, including removal of overhanging scrub and sediment, has proven highly effective in enhancing freshwater biodiversity. However, the implications of this management for farmland bird assemblages are unknown.

Bird surveys were undertaken at recently managed, open, macrophyte-dominated and at highly terrestrialised, macrophyte-free ponds in the intensively cultivated farmland of North Norfolk, UK. The diversity, abundance and composition of bird assemblages visiting these ponds were compared to determine responses to pond management by tree and mud removal.

Avian species richness, abundance and bird-visit frequencies were all higher at open farmland ponds. The observed patterns of bird occurrence were best explained by management-induced reductions in tree shading that resulted in aquatic macrophyte-dominance likely associated with high emergent invertebrate prey abundance. Moreover, we predict that open-canopy ponds offer greater habitat heterogeneity than overgrown ponds, allowing diversified bird use. Overgrown, terrestrialised ponds were preferred by some woodland bird species. Gamma diversity across the entire pondscape exceeded all individual pond alpha diversity measures by an order of magnitude, suggesting distinct variation in the bird assemblages visiting farmland ponds during different successional stages.

Pond management that generates a mosaic of pond successional stages, including open-canopy, macrophyte-dominated ponds, could help to address the long-term decline of farmland birds. We strongly advocate increased agro-ecological research in this field, combined with greater emphasis on ponds and pond management options in agri-environment schemes.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Landscapes in many parts of the world are dominated by farmland (Foley et al., 2005; Scherr and McNeely, 2008). Accordingly, agricultural landscapes have attracted substantial attention from the conservation research community. Historically, agricultural landscapes represented a highly dynamic habitat mosaic characterized by substantial spatio-temporal variations in environmental conditions (Chamberlain et al., 2000; Bennett et al., 2006). The resulting heterogeneity, at both local and regional scales, has been recognised as a primary factor underpinning historical agricultural landscape biodiversity (Benton et al., 2003; Tscharntke et al., 2005; Fahrig et al., 2011). Accordingly, increases

in agricultural intensification and associated agricultural habitat homogenization from the 1940s onwards, in combination with encroachments on remaining non-agricultural habitats, have resulted in a marked biodiversity reduction across the European countryside (Fuller, 2000; Ford et al., 2001; Robinson and Sutherland, 2002; Burel et al., 2004; Stoate et al., 2009; van Zanten et al., 2014).

Nearly 120 European bird species of conservation concern use lowland farmland habitats as either breeding or wintering habitat. A number of conservation priority species like the song thrush *Turdus merula*, yellowhammer *Emberiza citrinella* and reed bunting *Emberiza schoeniclus*, additionally rely on non-crop structures such as meadows, scrubland, woodlands, hedgerows and individual trees in agricultural landscapes as foraging, breeding and nesting sites (Whittingham et al., 2009; Marja and Herzon, 2012). Other birds such as skylark *Alauda arvensis* and grey partridge *Perdix perdix* are strongly affected by the quality of cropped habitats and marginal

* Corresponding author.

E-mail address: sarah.r.davies90@gmail.com (S.R. Davies).

habitats such as fallows and rough ground. Some 83% of European farmland bird species have undergone declines in abundance between 1970 and 1990 as a result of agricultural intensification. For 86% of these species, reductions were significant, and these trends have continued into the 21st century (Fuller et al., 1995; Donald et al., 2001; Barker, 2004; Holland, 2004; Butler et al., 2007; Baillie et al., 2014). Threats identified as affecting conservation priority bird species include the loss of old hedgerows, permanent pasture and scrub on farmland, changing sowing regimes, loss of variation in grassland swards, declines in abundance and diversity of insect prey, and reductions in seed resources linked to land-use changes and pesticide use (Chamberlain et al., 2000; Hinsley and Bellamy, 2000; Perkins et al., 2000; Donald et al., 2001; Benton et al., 2002; Barker, 2004; Holland, 2004).

While a range of approaches to enhance the farmed environment for wildlife have been taken in the UK and across Western Europe, many bird species populations have failed to recover (Donald and Evans, 2006). Declining UK Biodiversity Action Plan (BAP) species (JNCC, 2007) include skylark, starling *Sturnus vulgaris*, grey partridge and yellow wagtail *Motacilla flava* (Eaton et al., 2013). Aerial insectivorous birds associated with agricultural environments, such as swift *Apus apus* and house martin *Delichon urbicum*, have also shown steep population declines across industrialised European countries (Benton et al., 2002; Rioux Paquette et al., 2014). With farmland bird declines surpassing those in all other environments, serious concerns amongst both the scientific community and the general public have been raised. Currently, the main approach for counteracting farmland bird declines in Europe is the widespread adoption of agri-environment schemes (AES), such as the English Countryside Stewardship Schemes, but these have afforded limited success thus far for agricultural biodiversity (Kleijn et al., 2006, 2011; Baker et al., 2012).

A number of studies have concluded that agricultural management approaches that increase the heterogeneity of the agricultural mosaic will enhance overall species richness across many taxonomic groups at the landscape scale, while simultaneously improving ecosystem services and minimising agricultural yield losses (Pino et al., 2000; Auri and de Lucio, 2001; Weibull et al., 2003; Doxa et al., 2010; Sabatier et al., 2014). Soininen et al. (2015) stressed the importance of aquatic habitats for conservation, not only for aquatic organisms, but also for terrestrial species due to the contribution of potential cross-system subsidies from freshwater ecosystems which enhance terrestrial ecosystem functioning. Small wetlands, and especially ponds, may therefore play a crucial role in improving both aquatic and terrestrial biodiversity at the landscape scale, while also serving to increase habitat heterogeneity (Williams et al., 2004; Davies et al., 2008; C  r  ghino et al., 2008; Lemmens et al., 2013).

Ponds are of particular significance to biodiversity conservation in agricultural landscapes, forming habitat islands for a wide range of aquatic and semi-aquatic organisms in an otherwise species-poor environment (Declerck et al., 2006; Davies et al., 2008; Ruggiero et al., 2008). Unfortunately, many farmland ponds are threatened by in-filling (via land reclamation) and pollution due to agricultural intensification (Wood et al., 2003; Biggs et al., 2005; C  r  ghino et al., 2014). In addition, as a consequence of the general cessation of traditional pond management practices over the last 30–40 years (Sayer et al., 2013), a high proportion of UK farmland ponds have undergone terrestrialization, with the accumulation of litter and other organic material over time resulting in a decrease in pond size and depth. Many ponds also become increasingly encroached by woody vegetation and eventually transform into wet woodland, while in the absence of shrub and tree encroachment, pond succession can lead towards fen-swamp habitats. Indeed, in many areas, overgrown, tree-shaded ponds are

overwhelmingly dominant, resulting in sharp declines in landscape-scale aquatic diversity (Sayer et al., 2011, 2012). Approaches to combat widespread terrestrialisation include the creation of new ponds through initiatives such as the UK Million Ponds Project (Williams et al., 2010). As an alternative, existing, overgrown farmland ponds can be managed and restored via the removal of encroaching trees, scrub and accumulated pond sediment. The latter process effectively ‘resets’ succession thereby increasing the quality and quantity of open water habitats. Sayer et al. (2012) determined that macrophyte and invertebrate diversity was greatly enhanced in a managed pondscape comprising a mosaic of ponds at different successional stages set in an intensively managed agricultural landscape. Diversity patterns were strongly driven by degree of shading, with agricultural ponds previously deficient in macrophytes becoming macrophyte-dominated after management, providing habitat for a diverse array of species. Currently, both the UK Countryside Stewardship Scheme (CS) and Glacir Land Management Scheme for Wales offer options for maintaining and buffering ponds on farmland (Welsh Government, 2015; Natural England, 2015). Nonetheless, pond management itself is only included as a higher tier option within CS, and overall pond management remains relatively poorly promoted within UK AES.

While the influence of pond management on aquatic species assemblages is now established (Gee et al., 1997; Sayer et al., 2012), the links between pond management and the terrestrial environment have been comparatively neglected. Farmland ponds generally harbour substantial numbers of aquatic macroinvertebrates whose adult aerial stages are known to constitute an important food resource for nesting and fledging birds (Newton 1998; Baxter et al., 2005; Richardson et al., 2010; Schummer et al., 2012; Stenroth et al., 2015), and wintering waterbirds (Matuszak et al., 2014). In addition, mixed grassland margins around open ponds may increase the availability and diversity of broad-leaved plants and seeds utilised as a food resource by granivores (McCracken and Tallwin, 2004); we believe that these open pond margins are of high importance to birds.

We examine the value of a set of open, managed ponds and overgrown, non-managed ponds for bird communities in the intensively farmed agricultural landscape of North Norfolk, Eastern England. We predict that the benefits of pond management will strongly affect terrestrial organisms, as exemplified by the farmland bird community. The term ‘farmland bird’ in this context is used to encompass any species encountered within the agricultural landscape. This includes waterfowl, reed-nesting species, ground-nesting species and birds of prey, as well as open-country, woodland, scrubland and grassland bird species. We hypothesize that managed, macrophyte-dominated ponds attract a greater diversity of bird species than unmanaged, overgrown ponds, since they not only provide a higher diversity and abundance of emerging invertebrates and greater seed provision subsidy, but also increase habitat heterogeneity in the farmland landscape through provision of vegetated water and wet reed/sedge-dominated margins. We furthermore hypothesize that overgrown ponds primarily act as woodland habitat islands, occupied predominantly by woodland bird species. We finally hypothesize that bird assemblages use open and overgrown ponds for different activities in accordance with variations in habitat preference and food availability.

2. Methods

2.1. Study site

This study was conducted at four adjacent, intensive, mixed arable and cattle farms located between the villages of Melton




Fig. 1. Map of the farmland study area in Norfolk, eastern England, highlighting the open, managed ponds (W prefix) and overgrown ponds (D, M, S prefix) included in the study.

Constable, Stody and Briston in North Norfolk (Fig. 1). Most ponds in this region were created either for marl extraction or livestock watering between the 17th and 19th centuries (Prince, 1964). Of the 60 small ponds (<20 in diameter) in the 10 km² study area, a total of 22 ponds on privately owned farmland were selected for this study, thus allowing us to cover 36.6% of the pondscape. Selected ponds included 11 open canopy ponds with generally high submerged and fringing aquatic macrophyte cover (Fig. 2a,b) and 11 closed-canopy, overgrown ponds dominated by living and fallen trees of *Prunus spinosa*, *Salix* spp. and *Alnus glutinosa*, where aquatic plants were largely absent (Fig. 2c). All the ponds located in arable fields were surrounded by grassland buffers of at least 7 m width installed as part of Higher Level Stewardship (HLS) agreements.

The open canopy ponds were located at Manor Farm, Briston. Since the 1960s, most ponds at Manor Farm have been subject to a pond management programme comprising periodic scrub and pond sediment removal undertaken at two to four ponds each year with the aim of arresting terrestrialisation. This approach has created a mosaic of ponds varying in terms of degree of scrub encroachment and macrophyte cover. The resulting managed pondscape at Manor Farm is host to species-rich aquatic communities that include at least 16 breeding dragonfly species and the threatened Great Crested Newt, *Triturus cristatus*, which breeds in around 28 of the 40 ponds (Sayer et al., 2012, 2013). Moreover, the pondscape supports species-rich (n=24) communities of aquatic plants with frequent dominance of *Potamogeton*




Fig. 2. An open, managed pond (pond W10 in Fig. 1) at Manor Farm (a); swifts (*Apus apus*) feeding over pond W10 in May 2015 after a hatch of mayflies (b); a typical overgrown, highly terrestrialised pond in the study area (c).

natans in open water and fringing emergent vegetation typically including *Sparganium erectum*, *Typha latifolia* and *Epilobium hirsutum*.

2.2. Field surveys

All ponds included in the study were assigned an individual code depending on their location (Manor Farm ponds – W, Stody

ponds – S, Daniel's Farm ponds – D, Melton Constable ponds – M) (Fig. 1). Bird surveys were carried out at 11 managed, open ponds at Manor Farm, (W1–W37) and 11 late succession, overgrown ponds on surrounding farmland in the Stody/Hunworth/Melton Constable area (M1, D6, S4–S15). Most ponds formed isolated habitat “islands” on arable cropland or within cattle pastures where livestock were present at the time of the study (W1, W34, W37, S15). A number of ponds were along field boundaries and connected to hedgerow corridors (W34, W37, S4, S13, S15). On two occasions, ponds were situated within fields adjacent to patches of woodland (W1, M1). Three connected open-canopy ponds on Manor Farm were included in this study as a single pond complex, and thus are treated as one pond, termed the W22, 23, 24 cluster.

In June 2014, each of the 22 ponds was visited on five separate occasions, resulting in a total of 110 pond visits. Surveys were conducted during early mornings (5–10 am) and in good weather conditions, to avoid bias from lowered bird activity during periods of wet weather. Visit order was also randomized to avoid survey bias relating to time of day. During each individual pond visit, all birds encountered by sight, song or call were recorded over a set period of 20 min. Additional information recorded included the location of each bird individual on or around the pond (e.g. open water, surrounding vegetation, grassland buffer, stands of aquatic macrophytes) and bird behavioural activities: foraging, travelling, sheltering, vocal display, territorial behaviour, group behaviour and parental behaviour (including the provisioning of chicks). The activities and location of bird species for both pond types were subsequently compared by independent samples *t*-tests to determine differences in behaviour and habitat choice at open and overgrown ponds. Surveys were conducted at a location maximizing the visibility of the open pond surface area and the surrounding vegetation while minimising disturbance (Bibby et al., 1992). Active searches were also carried out around the circumference of each pond so that particularly large or obstructed sites could be viewed from different angles. All individual birds observed in or around the pond (up to 10 m away), including at directly adjacent trees, shrub and surrounding grassland buffer strips were recorded. Birds flying above the pond were also included, provided that they showed aerial feeding behaviour or flew low over the open water/tree canopy. Where possible, all encountered birds were identified to species level and subsequently grouped into ‘guilds’ based on avian family, diet (granivorous, insectivorous) and habitat preference (open country, scrubland, woodland, wetland, ground-nesting, reed nesting). In some instances, where sightings were very brief, distinguishing similar, closely-related species resulted in a high risk of misidentification. Species affected were the warbler genera *Phylloscopus* (chiffchaff/willow warbler), *Sylvia* spp. (garden warbler/blackcap), *Anas* spp. (mallard/gadwall) and *Motacilla* spp. (grey/yellow wagtail). Due to habitat preferences and species abundances, it was assumed that the respective unidentified female wagtails were yellow wagtails and female ducks were mallards, whereas in the other two cases, we combined all counts for the sets of two species and treated them as “super-species” in the statistical analysis.

Environmental data for each pond, including pond circumference, % pond surface shaded by trees, % pond circumference covered by trees, % coverage of pond surface by emergent (fringing) macrophytes and % coverage by submerged/floating-leaved macrophytes (assessed visually) were collected in 2012 and 2014. All aquatic plants were recorded on the DAFOR scale (Dominant – 5, Abundant – 4, Frequent – 3, Occasional – 2, Rare – 1) as described by Palmer et al. (1992), via visual assessments assisted by collections made using a double-headed rake.

2.3. Data analysis

Species richness, abundance and Simpson's and Shannon's diversity (Crist, 2003) were used to represent α - and γ -diversity, calculated by combining the records of the five individual pond visits. The highest recorded abundance for each bird species from all of the five surveys was used to represent the maximum number of individuals or “abundance” for each pond. Although this approach may still produce an overestimate of total bird abundance, the risk of counting the same individual multiple times is greatly diminished (Toms, 2004; BTO, 2014). Pond categories were subsequently compared using independent samples *t*-tests. Bird counts were rarefied using Hurlbert Rarefaction (Hurlbert, 1971) to create species accumulation curves for open and overgrown ponds. Correspondence Analysis (CA) was used to examine variation in bird assemblage composition between the ponds and to determine degree of species turnover between ponds (beta diversity) by maximizing the correspondence between species abundance scores and sample scores and measuring how distinct the sampling units were along gradients. Canonical Correspondence Analysis (CCA) was conducted to examine the direct relationships between pond environmental parameters and bird assemblages, again using bird abundance data. In addition, Stepwise Multiple Linear Regression (MLR) was performed to determine the extent to which environmental parameters were linked to overall bird diversity and abundance. Z-transformed environmental data were used in the multivariate analyses. Pearson's correlation analysis showed strong inter-correlations between four of the environmental parameters: submerged/floating and emergent macrophyte cover measures, % shading and % of the pond circumference surrounded by trees ($r \sim 0.7$). Pond circumference was an exception, however, and significantly correlated with percentage shading only ($r = -0.42$). Using the results of the Pearson's correlations a *p*-value threshold for parameter deletion of $p > 0.05$ was used, as values larger than this indicated that the effects of the variables upon patterns of avian diversity could not be separated. Subsequently, circumference and submerged/floating macrophyte coverage were chosen for further analysis, while the remaining parameters were omitted from MLR and CCA. It should be noted, however, that a high degree of submerged/floating macrophyte cover can be seen as a powerful proxy for low shading due to the highly negative correlation between these factors (Pearson's Correlation Coefficient, $r = -0.86$). Estimates *S* 8.2 was used in the calculation of both α - and γ -diversity (Colwell, 2009), while rarefaction curves were calculated using Species Diversity and Richness 3.02 (Pisces Conservation Ltd, 2002). CANOCO for Windows 4.5 (ter Braak and Smilauer, 2002) was used to generate CA and CCA ordination plots, while *t*-tests, stepwise MLR, and Pearson's Correlation Coefficient were all calculated in SPSS for Windows 20 (IBM Corp, 2011) and R: A Language and Environment for Statistical Computing Version 3.2.2 (R Core Team, 2015).

3. Results

3.1. Bird observations

In total, 58 breeding bird species were observed visiting or holding territories around the 22 farmland ponds (see Appendix A). Some 28 bird species were exclusive to only one pond type, while large proportions of the species encountered at open-canopy and overgrown ponds showed a very clear affinity to one of these pond type, as reflected by higher visit frequencies and abundances. Waterfowl, reed-associated species and open country species (comprising ground-nesting species, insectivorous open

country species and a number of granivorous species) were largely confined to open ponds. Nine of the eleven conservation priority UK BAP species that were recorded at the ponds showed a preference for open rather than overgrown ponds. Overall, the open ponds harboured a much higher diversity of bird species and guilds than overgrown ponds. Nonetheless, typical woodland bird species like the great spotted woodpecker *Dendrocopos major*, treecreeper *Certhia familiaris*, or nuthatch *Sitta europaea*, were exclusively encountered at overgrown ponds. Aside from 10 woodland bird species, all species recorded at overgrown ponds were also found at open ponds.

3.2. Avian diversity

Two of the largest, open-canopy farmland ponds, W10 and the W22/23/24 cluster, harboured the highest avian diversity (Species Richness, Abundance, Shannon's diversity, Simpson's Diversity); while the bird assemblage recorded at overgrown pond S7 was least diverse. However, Shannon's and Simpson's Diversity did not differ significantly between individual ponds of each type (Table 1). Nonetheless, both the recorded and estimated species richness was significantly higher at open ponds compared to overgrown ponds ($p < 0.05$, Table 1). This trend was further supported by rarefaction curves combining samples of the two groups (Fig. 3). Gamma diversity across the agricultural pondscape was considerably higher than both the alpha diversity of any one individual pond, and of the combined open and overgrown ponds, indicating important diversity contributions by both pond types (Table 1).

In the CA bi-plot, axis 1 explained 15.3% of species data variance, whereas axis 2 explained a further 10.5%. Species turnover between the overgrown ponds was relatively low, as illustrated by the small area of ordination space generally occupied by these sites in the CA (Pond S7 is an outlier due to a record of tawny owl *Strix aluco*, Fig. 4). In contrast, a greater bird species turnover was observed at the open ponds, meaning that these are more heterogeneous in the bird assemblages they support. The bird community structure showed significant variation in relation to the measured environmental gradients in the agricultural pondscape. In the CCA bi-plot (Fig. 5), axis 1 was positively related to macrophyte coverage (and thus negatively correlated with shading) and explained 9.99% of bird species' variance. Axis 2, which explained an additional 4.43% of bird species' variance, was strongly associated with pond circumference. Bird species were widely distributed across axis 1, showing varying preferences for macrophyte coverage and associated shading, but generally the species most prevalent at open ponds, such as aerial insectivores (swift, swallow, house martin), open country species (e.g. whitethroat *Sylvia communis*, linnet *Carduelis cannabina*), granivores (e.g. greenfinch *Chloris*, skylark, house sparrow *Passer*




Fig. 3. Hurlbert rarefaction curves for overgrown ponds, open ponds and all ponds combined. Number of individuals sampled plotted against number of species encountered with error bars representing the standard error (\pm SE).

domesticus), dabbling ducks (mallard *Anas platyrhynchos* and gadwall *Anas strepera*) and wetland passerines (reed bunting, reed warbler *Acrocephalus scirpaceus* and sedge warbler *Acrocephalus schoenicus*), preferred lower levels of tree shading (and thus high macrophyte coverage), whereas woodland birds such as robin *Erithacus rubecula*, nuthatch and treecreeper were associated with increased tree shading (and thus lower macrophyte coverage). Birds associated with ponds of intermediate macrophyte coverage and partial shading (e.g. coal tit, long-tailed tit), as well as species equally abundant at both open and overgrown ponds (e.g. chaffinch, yellowhammer, blue tit), clustered towards the centre of the plot.

MLR indicated that pond circumference and macrophyte coverage were both significant predictors of overall avian species richness ($F = 11.82$, Adjusted $R^2 = 0.51$, $p = 0.0004$), and abundance ($F = 12.32$, Adjusted $R^2 = 0.52$, $p = 0.0003$) (Table 2). While circumference was a significant predictor for Shannon's Diversity (estimate = 0.04, t value = 2.74, $p = 0.01$), this was not true for macrophyte coverage (estimate = 0.35, t value = 1.22, $p = 0.35$). In addition, the model failed to explain the patterns in Simpson's Diversity ($p = 0.95$).

3.3. Pond use by farmland birds

In addition to vegetated open water, the open agricultural ponds afforded a variety of associated habitats that were utilised by birds, and a number of bird behaviours were observed more

Table 1

Diversity and abundance measures comparing avian alpha diversity of open and overgrown ponds and gamma diversity of birds from all ponds, where figures for alpha diversity measures represent mean values \pm standard error of the mean.

Pond category	Species Richness ($\bar{x} \pm \text{SE}$)	Abundance (no. individuals) ($\bar{x} \pm \text{SE}$)	Shannon's Diversity ($\bar{x} \pm \text{SE}$)	Simpson's Diversity ($\bar{x} \pm \text{SE}$)
Alpha Diversity				
Open	17.5 \pm 1.4 [*]	38.3 \pm 3.8 ^a	13.7 \pm 0.9 ^b	16.6 \pm 1.5 ^b
Overgrown	13.3 \pm 0.7 [*]	24.4 \pm 1.8 ^a	11.5 \pm 0.7 ^b	18.2 \pm 1.8 ^b
Gamma Diversity				
All Ponds	58	679	31.4	24.1
Combined Open	46	421	28.8	23.2
Combined Overgrown	35	268	21.1	17.6

Statistical significance of independent samples t -tests comparing alpha diversity means of open and overgrown ponds are based on the p -value threshold of $p < 0.05$ and are denoted by ^{a,b}.

^{*} $p = 0.02$.

^a $p = 0.00072$.

^b $p > 0.05$.




Fig. 4. Correspondence Analysis (CA) of pond site (a) and bird species (b) data. Ponds are coded according to treatment (open, managed or overgrown, unmanaged).

Key to species codes: BARO: barn owl, BLAB: blackbird, BLAC: blackcap, BLAG: black-headed gull, BLUT: blue tit, BULF: bullfinch, CARC: carrion crow, CHAF: chaffinch, CHIC: chiffchaff, COAT: coal tit, COLD: collared dove, DUNN: dunnock, GADW: gadwall, GARW: garden warbler, GOLC: goldcrest, GOLF: goldfinch, GREF: greenfinch, GREP: grey partridge, GRET: great tit, GRSW: great-spotted woodpecker, HOUM: house martin, HOUS: house sparrow, JACD: jackdaw, JAY: jay, LINN: linnet, LWHT: lesser whitethroat, LONT: long-tailed tit, MAGP: magpie, MALL: mallard, MART: marsh tit, MEAP: meadow pipit, MIST: mistle thrush, MOOH: moorhen, NUTH: nuthatch, NIGH: nightingale, OYSC: oystercatcher, PHEA: pheasant, PIEW: pied wagtail, REDP: red-legged partridge, REDS: redstart, REEB: reed bunting, REEW: reed warbler, ROBI: robin, SEDW: sedge warbler, SKYL: skylark, SPAH: sparrowhawk, SPOF: spotted flycatcher, SONT: song thrush, STAR: starling, STOC: stonechat, SWAL: swallow, SWIF: swift, SYLV: Sylvia/unidentified warbler (genus), TAWO: tawny owl, TREC: treecreeper, WHIT: whitethroat, WILW: willow warbler, WOOP: wood pigeon, WREN: wren, YELH: yellowhammer, YELW: yellow wagtail.

frequently at open ponds (Table 3, Appendix B Tables B1 and B2). At the open ponds, foraging was a particularly important activity, and was significantly more prevalent at open ponds compared to overgrown ponds ($t = 2.44$, $df = 10$, $p = 0.03$), especially amongst open country, insectivorous species such as swallows, swifts and whitethroats. Further, many open-country bird species such as linnet, yellowhammer, reed bunting, house sparrow and greenfinch, as well as the ground-nesting grey partridge and skylark, were strongly associated with the grassland buffer strips around open ponds, but did not show a similar affinity to buffer strips at overgrown ponds ($t = 2.97$, $df = 10$, $p = 0.01$). Emergent plant stands

(e.g. sedge beds), which were utilised at open ponds by ducks (*Anas* spp.) and warblers (*Acrocephalus* spp.) were furthermore widely lacking at overgrown ponds, leading to an associated absence of these species. Tree vegetation at both pond groups was important for refuge and as a perch for singing and territorial displays ($t = -0.61$, $df = 10$, $p = 0.55$). However, breeding pairs and family groups, occasionally even nesting within the pond cluster, were observed more frequently at open ponds ($t = 3.74$, $df = 10$, $p = 0.003$), and evidence of chick provisioning was recorded on more occasions at open ponds ($t = 2.5$, $df = 10$, $p = 0.03$). Aside from a few aquatic species such as moorhen *Gallinula chloropus*, the bird species encountered at overgrown ponds were largely confined to the surrounding wet woodland vegetation rather than the water-body itself.

4. Discussion

4.1. Drivers of avian diversity at farmland ponds

Similar to previous studies (Froneman et al., 2001; Sebastián-González et al., 2010), larger ponds possessed a larger pool of bird species. The most species-rich ponds however were not only large, but also harboured abundant and spatially heterogeneous macrophyte communities. Macrophytes are extremely important components of pond ecosystems, with high macrophyte coverage exerting a significant positive influence on overall aquatic diversity (McAbendroth et al., 2005; Thomaz and da Cunha, 2010; Florencio et al., 2014). Generally, the influence of open-canopy, macrophyte-dominated ponds on both aquatic and terrestrial species has to date largely evaded scientific research. Our results show that the abundance and diversity (species richness) of birds encountered in the direct vicinity of ponds was strongly positively influenced by macrophyte coverage, and strongly negatively associated with high levels of shading, although it is difficult to identify underlying causal relationships.

Under conditions of high tree/scrub shading at late-successional unmanaged ponds, aquatic plants are typically eliminated. By contrast, management-induced reductions in shading lead to a rapid, positive response of aquatic macrophytes in terms of both species cover and diversity (Sayer et al., 2012). Presence of vegetation within ponds is cited as an important factor for waterbirds when selecting wetland habitat (Cody 1985; Sebastián-González et al., 2010), since increased macrophyte cover provides benefits such as food, nesting material, habitat and refuge from predators (McKinstry and Anderson, 2002; Santoul et al., 2009). Our results show that such benefits extend beyond water-birds to birds encountered across agricultural landscapes more generally, covering open country, ground nesting, reed nesting, granivorous and insectivorous guilds, all of which appeared to associate with, and potentially benefit from, open-canopy, macrophyte-dominated ponds and their connected grassland buffers. Notably, open-canopy ponds appeared to offer suitable habitat for a number of UK BAP conservation-priority farmland species and species undergoing declines on farmland, such as skylark, grey partridge and reed bunting, all of which were primarily associated with open ponds, but were absent at overgrown ponds. The habitat associations of these species suggest that individuals can find some of the nesting or foraging resources required for their persistence in or around open ponds.

4.2. Pond habitat and food resources for birds

The higher richness and abundance of bird species using open-canopy ponds could be the result of a variety of ecological mechanisms, particularly those relating to habitat complexity, the high degree of habitat variation among individual managed ponds




Fig. 5. Canonical Correspondence Analysis (CCA) showing pond site (a) and species (b) data. Ponds are coded according to treatment (open, managed or overgrown, unmanaged).

Key to species codes: BARO: barn owl, BLAB: blackbird, BLAC: blackcap, BLAG: black-headed gull, BLUT: blue tit, BULF: bullfinch, CARC: carrion crow, CHAF: chaffinch, CHIC: chiffchaff, COAT: coal tit, COLD: collared dove, DUNN: dunnoek, GADW: gadwall, GARW: garden warbler, GOLC: goldcrest, GOLP: goldfinch, GREP: greenfinch, GREP: grey partridge, GRET: great tit, GRSW: great-spotted woodpecker, HOUM: house martin, HOUS: house sparrow, JACD: jackdaw, JAY: jay, LINN: linnet, LWHT: lesser whitethroat, LONT: long-tailed tit, MAGP: magpie, MALL: mallard, MART: marsh tit, MEAP: meadow pipit, MIST: mistle thrush, MOOH: moorhen, NUTH: nuthatch, NIGH: nightingale, OYSC: oystercatcher, PHEA: pheasant, PIEW: pied wagtail, REDP: red-legged partridge, REDS: redstart, REEB: reed bunting, REEW: reed warbler, ROBI: robin, SEDW: sedge warbler, SKYL: skylark, SPAH: sparrowhawk, SPOF: spotted flycatcher, SONT: song thrush, STAR: starling, STOC: stonechat, SWAL: swallow, SWIF: swift, SYLV: Sylvia/unidentified warbler (genus), TAWO: tawny owl, TREC: tree creeper, WHIT: whitethroat, WILW: willow warbler, WOOP: wood pigeon, WREN: wren, YELH: yellowhammer, YELW: yellow wagtail.

and increased food availability. Aquatic invertebrates are known to establish much more diverse communities in structurally-complex macrophyte stands associated with open ponds, which results in both a greater diversity and abundance of adult stages (Gee et al., 1997; McAbendroth et al., 2005; Hinden et al., 2005), and following emergence and dispersal from the pond may form an important food subsidy for foraging insectivorous birds (Schummer et al., 2012; Dreyer et al., 2015; Fig. 6). Key potential invertebrate prey taxa include the orders Odonata, Ephemeroptera and Coleoptera, and the family Chironomidae. In our study, we did not quantify

emergent invertebrate abundance or diversity; however, our previous research showed that, with the exception of molluscs, managed Manor Farm ponds showed higher invertebrate diversity than unmanaged ponds, with invertebrate diversity steadily increasing for 3–5 years after management (Sayer et al., 2012). In our present study, observations at open ponds suggested that adult invertebrate prey were abundant. Aerial insectivores such as swallows, swifts and house martins seemed primarily driven by emerging invertebrates, and pairs or groups were frequently observed hovering, diving and catching insects on the wing over open water (as in Fig. 2b). Invertebrate resources offered by ponds may become particularly important during the breeding season, when nutritional requirements are elevated. A number of whitethroat nests were encountered in the bushes fringing open canopy ponds and adults were regularly observed provisioning young. Nesting sites adjacent to open ponds may have been favoured by this species to allow better access to invertebrate-rich foraging sites when provisioning offspring.

A variety of grasses, sedges, rushes and herbs of different heights and structures were encountered around the open ponds (Fig. 2a,b), which may offer nesting materials, seed resources, refuge from predators and resting and perching habitat, as well as important habitat for invertebrate prey. Josefsson et al. (2013) observed that fields with grassland buffer strips supported significantly more skylark territories than fields without buffer strips, with such sites characterized by increased densities of spiders and beetles. Thus, for farmland birds that rely on the cropped area of fields for both breeding and foraging (such as skylark), grassland buffer strips around isolated, open farmland ponds could play an important supplementary role in terms of food resources, provided that there is a sufficient density of ponds in the landscape. In a pondscape setting, we suggest that surrounding grassland margins may act as recipients of particularly high numbers of invertebrate prey originating from the pond, with invertebrate assemblages in these buffers further enhanced by the presence of humidity gradients from the pond margin towards agricultural habitats on higher ground (Fig. 6a). Seeds associated with pond marginal areas may also form an important part of the diet of many conservation priority granivores on farmland, including house sparrow, yellowhammer and linnet (Atkinson et al., 2004; Robinson et al., 2004; McCracken and Tallowin, 2004). A key, known bottleneck for farmland birds is starvation in late winter, a phenomenon known as the “winter hungry gap” (Siriwardena et al., 2008). It is possible that particularly open, plant-rich ponds may provide a seed-rich area that persists through winter and thus assists bird survival. In contrast, the grass and dicot seeds involved would not be available in overgrown habitats because both the plants and birds concerned are open country species: the plants are not found in shaded conditions and the birds forage in open areas, not within woody vegetation.

The lower species diversity observed at overgrown ponds is probably due to the relative homogeneity of habitats offered by such ponds, which essentially mimic small wet woodland sites. Although the overgrown ponds were also surrounded by grassland buffers, these apparently failed to offer birds the same benefits as grassland buffers around open ponds, possibly because the grassland was heavily shaded and separated from the pond by a dense barrier of woody vegetation (Fig. 2c,b). Open country species often avoid vertical structures (Sparks et al., 1996), rendering areas immediately around densely wooded ponds unattractive to these species. It follows that another possible cause of lower avian diversity at overgrown ponds is a perceived heightened risk of ambush from predators around dense cover, particularly for open country species (Cresswell, 1996). Although it will not have represented a real predatory threat, a tawny owl observed at overgrown pond S7 may have affected what was detected there:

Table 2
Stepwise Multiple Linear Regression (MLR) showing results for effects of (i) Submerged/floating macrophyte coverage and (ii) Circumference on test variables (a) Species richness, (b) Abundance and (c) Shannon's diversity.

Test Variable and Predictors	Beta Coefficient	t-value	p-value
a) Species richness			
i) Submerged/floating macrophyte coverage	0.74	2.15	0.044*
ii) Circumference	0.06	3.51	0.0002*
Adjusted $R^2 = 0.51$, $F_{(2,19)} = 11.82$, $p = 0.0004^*$			
b) Abundance			
i) Submerged/floating macrophyte coverage	2.31	2.32	0.03*
ii) Circumference	0.18	3.51	0.002*
Adjusted $R^2 = 0.52$, $F_{(2,19)} = 12.32$, $p = 0.0003^*$			
c) Shannon's diversity			
i) Submerged/floating macrophyte coverage	0.35	1.22	0.35
ii) Circumference	0.04	2.74	0.01*
Adjusted $R^2 = 0.32$, $F_{(2,19)} = 5.96$, $p = 0.009^*$			

* Statistical significance is based on the p -value threshold of $p < 0.05$.

Table 3
Frequencies of behaviours and locations of birds recorded at open and overgrown ponds. Values are given as means \pm standard error of the mean (SEM) of the 11 open ponds and 11 overgrown ponds, along with corresponding t -values from the independent samples t -tests.

	Behaviours observed		Recorded locations of birds		
	Foraging	Provisioning offspring	Pair/family groups	Grassland buffer	Tree vegetation
Open	15.6 \pm 4.1	4.18 \pm 1.4	25.5 \pm 4	6.45 \pm 2.2	33.8 \pm 6.2
Overgrown	5.36 \pm 1.3	0.63 \pm 0.3	8.27 \pm 1.6	0.1 \pm 0.1	38.7 \pm 2.7
t -value	2.44*	2.5*	3.74 ^a	2.97 ^a	-0.61

Statistical significance for independent samples t -tests is based on the p -value threshold of $p < 0.05$ and is denoted by *.^a

* $p < 0.05$.

^a $p < 0.01$.




Fig. 6. Conceptual diagrams depicting habitat features and resources for farmland birds at typical open managed ponds (a) and overgrown ponds (b).

this pond was distinct from the other overgrown ponds not least in supporting the lowest number of bird species.

It could be argued that a lack of bird diversity observed amongst the overgrown ponds was partly an artefact of reduced visibility at overgrown ponds. However, while birds may not always have been seen at these ponds, hidden birds still had a high chance of detection by their vocalisations. Clearly, overgrown ponds also afforded good habitat for woodland birds. In this respect, they may be used as stepping stones for species travelling between larger woodland sites (Neuschultz et al., 2013). Therefore, maintaining some overgrown ponds should have positive implications for

habitat connectivity, promoting the dispersal of woodland species (Lawton et al., 2010).

4.3. Pond management and farmland bird conservation

This study suggests that pond management can be considered to be a valuable tool for bird conservation in farmland. It also alludes to the importance of maintaining a mosaic of pond successional stages within agricultural landscapes in order to support a wide variety of bird guilds. However, the relative value of each successional stage will depend on the extent to which it

contributes to the existing habitat heterogeneity in a given landscape. The continued terrestrialisation of entire agricultural pondscape risks eliminating the contribution of open ponds to landscape-level avian diversity. Equally, simultaneous, uniform pond management with associated loss of wet woodland habitat and homogenisation of the pondscape could have detrimental effects for woodland guilds, particularly declining wet woodland species such as marsh tit, which was uniquely associated with semi-overgrown ponds in this study. We recommend that a high level of environmental variability should be maintained across agricultural pondscape, taking resource and habitat requirements of specialist bird groups most at risk from future declines into account (Gregory et al., 2004; Le Viol et al., 2012).

Clearly, the Manor Farm approach of arresting succession at just a few ponds every year, with some ponds left to natural development, ensures the existence of a pond mosaic comprising ponds of varying stages of succession, which could provide an ideal scenario for farmland bird conservation. In other regions, where ponds are less abundant, creation of new ponds could be required to provide new habitat for local bird populations. We predict that the benefits of pond management for biodiversity are by no means confined to the aquatic environment or even the immediate vicinity of the pond. Instead, where many open ponds are present, high rates of aquatic invertebrate deposition and dispersal may significantly increase invertebrate abundance and diversity across the entire landscape through a strong “chimney effect” (Fig. 6a). As aquatic and terrestrial ecosystems are tightly linked (Knight et al., 2005), increasing the interchange of resources between aquatic and terrestrial habitats might be of paramount significance to regional biodiversity (Baxter et al., 2005; Richardson et al., 2010), with cross-system subsidies, represented here by aquatic insect deposition and food-plant resources, being of significant importance to ecosystem functioning in farmland environments (Allen et al., 2012; Bartels et al., 2012; Dreyer et al., 2015; Soininen et al., 2015).

Further study is needed to quantify emergent invertebrate abundance and diversity at managed and un-managed ponds, as well as to determine how pond management may be optimized to enhance both breeding and overwintering of farmland birds. Our study is limited in its spatial and temporal coverage, and we suggest that future bird, macrophyte and invertebrate surveys are carried out at different times of the year to account for seasonal variability. This should lead to much improved understanding of the role of ponds for farmland bird conservation. Nevertheless, our study strongly suggests that pond management has a very important role to play in this respect. Ponds are cheap and simple to manage compared to other habitats, yet they remain a rarely promoted option within AES. We propose that more emphasis be placed on the value of ponds and their management within agricultural policy, environmental education and conservation strategies, within and across the farmed landscape.

Acknowledgements

We wish to thank all the landowners of the farms used in this study for permitting us to survey their ponds, E. Smith for assistance with field work and D. and J. Sayer for much logistical support. We also thank the UCL Environmental Change Research Centre for financial support, G. Hilton (WWT) and S. Newson (BTO) for advice on bird survey methods, and the UCL MSC in Conservation for general facilitation. We dedicate this study to R. Waddingham at Manor Farm, who came up with all the ideas that underpin this work. The dedication and care he demonstrates towards his ponds show that nature conservation and modern agriculture can coexist in a highly sustainable way – a powerful and much needed message.

Appendix A.

See Table A1.

Table A1

Species records for both open managed and overgrown, non-managed ponds. Species are grouped according to guild or habitat preference and sub-divided into families and allies.

Habitat/Guild	Species Name	Common Name	Open Ponds ^b	Overgrown Ponds ^b
Waterfowl and Rallids	<i>Anas platyrhynchos</i>	Mallard	W9(1), W16(3), W17(3), W19(2), W33(1), W22/23/24(1), W34(1)	
	<i>Anas strepera</i>	Gadwall	W16(2), W34(1)	
	<i>Gallinula chloropus</i>	Moorhen	W8(1), W9(4), W10(4), W16(3), W17(4), W19(3), W22/23/24(2), W33(3), W34(2), W37(3)	S5(1), D6(1), S9(4), S10(2), S11(3), S13(4)
	<i>Carduelis cannabina</i>	Linnet ^c	W10(1), W22/23/24(1), W33(2), W34(1)	
Seed Eating ^a Finches and Allies	<i>Carduelis carduelis</i>	Goldfinch	W1(1), W10(1), W16(3), W19(1), W22/23/24(3), W34(1), W37(3)	M(1), S4(2), D6(1), S9(1), S11(1), S13(1)
	<i>Chloris chloris</i>	Greenfinch	W8(1), W9(1), W16(1), W33(1),	
	<i>Emberiza citrinella</i>	Yellowhammer ^c	W8(3), W9(1), W10(2), W16(3), W17(4), W19(1), W22/23/24(2), W33(3), W37(1)	S4(3), S5(1), D6(1), S6(1), S7(3), S9(1), S10(1), S11(2), S13(1), S15(3)
	<i>Fringilla coelebs</i>	Chaffinch	W1(2), W8(4), W9(2), W10(5), W16(3), W17(4), W19(1), W22/23/24(4), W33(2), W34(3), W37(3)	M1(1), S4(1), S5(2), D6(4), S6(1), S7(4), S9(4), S10(4), S11(4), S13(2), S15(5)
	<i>Passer domesticus</i>	House Sparrow ^c	W23(2), W33(1) W34(2)	
	<i>Pyrrhula pyrrhula</i>	Bullfinch ^c	W22/23/24(1), W4(1)	
	<i>Emberiza schoeniclus</i>	Reed Bunting ^c	W16(1), W17(3), W23(1), W33(2), W34(1)	
Seed Eating ^a	<i>Alauda arvensis</i>	Skylark ^c	W1(1), W8(1), W9(1), W10(2), W16(4), W17(2), W19(1), W33(2)	
Grass Nesting	<i>Alectoris rufa</i>	Red-Legged Partridge	W17(1)	
	<i>Perdix perdix</i>	Grey Partridge ^c	W16(1)	
	<i>Phasianus colchicus</i>	Pheasant	W19(1)	
	<i>Dendrocopos major</i>	Great Spotted Woodpecker		S5(1), S6(1)
Insectivorous Woodland		Robin	W10(1), W22/23/24(2)	

Table A1 (Continued)

Habitat/Guild	Species Name	Common Name	Open Ponds ^b	Overgrown Ponds ^b
(Thrushes and Allies)	<i>Erithacus rubecula</i>			M1(2), S4(4), S5(4), D6(3), S6(1), S11(4), S10(5), S9(3), S15(1)
	<i>Luscinia megarhynchos</i>	Nightingale	W8(1), W22/23/24(1), W17(1)	
	<i>Phoenicurus phoenicurus</i>	Redstart		S10(2)
	<i>Turdus merula</i>	Blackbird	W1(2), W8(1), W22/23/24 (2), W33(1)	M1(3), S4(2), S5(2), D6(3), S9(2), S10(2), S13(2)
	<i>Turdus viscivorus</i>	Song Thrush ^c		S9(1), S10(1)
Insectivorous Woodland	<i>Aegithalos caudatus</i>	Mistle Thrush		S4(1), S15(1)
	<i>Cyanistes caeruleus</i>	Long Tailed Tit	W10(1), W22/23/24(2)	S11(1), S13(1)
	<i>Parus major</i>	Blue Tit	W(3) W8(1), W9(3), W10(3), W16(1), W17(1), W19(2), W22/23/24 (4), W34(1), W37(2)	M1(2), S4(5), S5(2), D6(2), S6(3), S7(3), S13(3), S9(2), S10(3), S11(5), S15(3)
(Paridae and Allies)	<i>Periparus ater</i>	Great Tit	W1(2), W8(1), W9(1), W10(3), W16(1), W17(1), W22/23/24(1), W34(2), W37(1)	S4(3), S5(1), S6(1), S9(2), S10(1), S11(2), S13(3), S15(2)
	<i>Poecile palustris</i>	Coal Tit	W1(1), W10(1) W22/23/24(1)	
	<i>Certhia familiaris</i>	Marsh Tit ^c	W1(1)	M1(2)
		Treecreeper		M1(1), S15(2)
	<i>Sitta europaea</i>	Nuthatch		S5(1), D6(1), S15(2)
Insectivorous Woodland	<i>Troglodytes troglodytes</i>	Wren	W1(1), W8(1), W9(2), W10(4), W19(2), W22/23/24(2), W34(4), W37(4)	M1(4), S4(1), S5(1), D6(3), S6(3), S7(5), S9(4), S10(4), S11(4), S13(5), S15(4)
	<i>Muscicapa striata</i>	Spotted Flycatcher ^c	W10(2), W22/23/24(1), W33(1) W34(1)	D6(1), S6(1), S9(1)
	<i>Phylloscopus collybita/trochilus</i>	Chiffchaff/Willow Warbler	W1(1), W10(4), W19(1), W22/23/24(1), W37(1)	M1(1), D6(1), S4(2), S7(1), S9(3), S10(3), S11(3), S13(2), S15(4),
	<i>Prunella modularis</i>	Dunnock	W10(1), W33(2), W34(1)	S4(1), S11(1), S13(1), S15(2)
(Warblers and Allies)	<i>Regulus regulus</i>	Goldcrest	W33(1), W34(1)	S11(1)
	<i>Sylvia borin/atricapilla (G)</i>	Garden Warbler/Blackcap	W8(2), W9(1), W10(1), W16(1), W17(1), W22/23/24(3), W33(1),	M1(1), D6(3), S4(1), S10(3), S11(1), S13(1)
	<i>Sylvia communis</i>	Whitethroat	W1(2), W9(2), W10(3), W16(1), W17(1), W19(1), W22/23/24(4), W33(4) W34(1)	D6(2), S7(3)
	<i>Sylvia curruca</i>	Lesser Whitethroat	W17(1), W22/23/24(4), W34(1), W37(1)	
	<i>Acrocephalus schoenobaenus</i>	Sedge Warbler	W9(3)	
Reed Nesting Warblers	<i>Acrocephalus scirpaceus</i>	Reed Warbler	W10(1)	
	<i>Anthus pratensis</i>	Meadow Pipit		M1(1), S4(1)
	<i>Motacilla alba</i>	Pied Wagtail		S5(1)
Insectivorous Open Country (Pipits and Wagtails)	<i>Motacilla flava</i>	Yellow Wagtail ^c	W33(1)	
	<i>Apus apus</i>	Swift	W10(3), W22/23/24(1), W33(1), W34(1)	S13(1)
	<i>Delichon urbicum</i>	House Martin	W10(1), W19(2), W33(1)	
Corvids	<i>Hirundo rustica</i>	Swallow	W16(2), W9(1), W10(3), W17(3), W33(3), W34(4), W37(1)	S13(1)
	<i>Corvus corone</i>	Carriion Crow	W9(1) W19(1)	M1(1), S10(1), S11(1), S13(1), S15(1)
	<i>Corvus monedula</i>	Jackdaw	W17(1), W19(1), W34(1)	S9(1), S13(2)
	<i>Garrulus glandarius</i>	Jay		D6(1)
Doves	<i>Pica pica</i>	Magpie	W10(2)	S5(1), S13(2)
	<i>Columba palumbus</i>	Woodpigeon	W1(3), W8(2), W9(3), W10(2), W16(1), W17(1), W19(1), W22/23/24(3), W34(2)	M1(1), S13(4), S5(5), D6(1), S6(2), S7(2), S9(2), S10(2), S11(2), S15(5)
	<i>Streptopelia decaocto</i>	Collared Dove	W1(1)	S9(1)
	<i>Accipiter nisus</i>	Sparrowhawk	W19(1)	
Birds of Prey	<i>Strix aluco</i>	Tawny Owl		S7(1)
	<i>Tyto alba</i>	Barn Owl	W1(1)	
Shorebirds	<i>Haematopus ostralegus</i>	Oystercatcher		S13(1)
	<i>Larus melanocephalus</i>	Black-Headed Gull	W19(1), W22/23/24(1), W33(2), W34(1)	S13(1)

W = Manor Farm ponds, S = Stody/Hunworth ponds, M = Melton Constable ponds, D = Daniel's ponds.

G Classification to genus level only.

^a Seed eaters which become insectivorous during the breeding season and when provisioning young.^b Values inside brackets indicate frequency of visits i.e. number of surveys present out of a total of five (scores = 1–5).^c Indicates UK Biodiversity Action Plan (BAP) species.

Appendix B.

Table B1

Observations of the activities and locations of 12 observed bird species at open, managed ponds.

^aFor foraging data, brackets indicate number of times individuals were observed with prey items in mouth.^bConfirmation of young accompanied by adults in the case of waterfowl and rallids.

Species Name Common Name	<i>Hirundo rustica</i> Swallow	<i>Alauda arvensis</i> Skylark	<i>Gallinula chloropus</i> Moorhen	<i>Anas platyrhynchos</i> Mallard	<i>Emberiza schoeniclus</i> Reed Bunting	<i>Emberiza citrinella</i> Yellowhammer	<i>Fringilla coelebs</i> Chaffinch	<i>Cyanistes caeruleus</i> Blue Tit	<i>Troglodytes troglodytes</i> Wren	<i>Phylloscopus collybita</i> Chiffchaff	<i>Sylvia communis</i> Whitethroat	<i>Turdus merula</i> Blackbird
Foraging/ hunting	39(1) ^a	0	0	4	0	0	5	19	0	0	20(5)	1(1)
Pair/group behaviour	35	9	8	16	4	2	29	36	0	0	16	0
Evidence of provisioning chicks	2	0	2 ^b	4 ^b	0	0	4	2	0	0	18	1
Sheltering/ using cover	1	11	31	8	8	15	35	37	23	4	34	5
Perching	1	5	4	0	9	14	44	53	24	4	29	5
Territorial behaviour; singing	2	16	0	0	2	18	21	28	19	4	12	2
Calling	15	0	24	5	3	1	7	14	4	0	21	4
Riparian vegetation; shrubs and trees	1	6	18	8	6	14	54	55	21	4	30	6
Reeds/rushes	0	1	17	1	2	0	0	0	0	0	7	0
Grassland buffer	0	8	0	0	5	3	2	3	3	0	9	0
Bank/pond edge	0	0	16	4	1	0	8	2	2	0	4	1
Swimming in open water	0	0	13	13	0	0	0	0	0	0	0	0

^{*}Evidence of provisioning chicks was determined by the occurrence of repeated visits to a site suspected to contain a nest or chicks by pairs of groups to the same site. Includes observations of individuals bringing food items to the site and/or taking turns to forage and guard territory.^aFor foraging data, brackets indicate number of times individuals were observed with prey items in mouth.^bConfirmation of young accompanied by adults in the case of waterfowl and rallids.**Table B2**

Observations of the activities and locations of 12 observed bird species at overgrown, terrestrialised ponds.

Species Name Common Name	<i>Hirundo rustica</i> Swallow	<i>Alauda arvensis</i> Skylark	<i>Gallinula chloropus</i> Moorhen	<i>Anas platyrhynchos</i> Mallard	<i>Emberiza schoeniclus</i> Reed Bunting	<i>Emberiza citrinella</i> Yellowhammer	<i>Fringilla coelebs</i> Chaffinch	<i>Cyanistes caeruleus</i> Blue Tit	<i>Troglodytes troglodytes</i> Wren	<i>Phylloscopus collybita</i> Chiffchaff	<i>Sylvia communis</i> Whitethroat	<i>Turdus merula</i> Blackbird
Foraging/ hunting	1	0	0	0	0	0	4	6	4	2	0	3(1) ^a
Pair/group behaviour	0	0	2	0	0	2	8	19	0	6	0	0
Evidence of provisioning chicks [*]	0	0	5 ^b	0	0	0	0	0	0	0	0	1
Sheltering/ using cover	0	0	14	0	0	19	29	39	41	21	2	16
Perching	1	0	0	0	0	19	40	52	31	24	5	18
Territorial behaviour; singing	0	0	0	0	0	20	23	34	35	21	5	3
Calling	0	0	15	0	0	0	6	32	15	3	0	12
Riparian vegetation; shrubs and trees	0	0	11	0	0	22	41	59	53	24	5	17
Reeds/rushes	0	0	14	0	0	0	0	0	3	1	0	0
Grassland buffer	0	0	0	0	0	0	0	0	0	0	0	0
Bank/pond edge	0	0	13	0	0	0	0	1	17	4	1	5
Swimming in open water	0	0	7	0	0	0	0	0	0	0	0	0

^{*}Evidence of provisioning chicks was determined by the occurrence of repeated visits to a site suspected to contain a nest or chicks by pairs of groups to the same site. Includes observations of individuals bringing food items to the site and/or taking turns to forage and guard territory.^aFor foraging data, brackets indicate number of times individuals were observed with prey items in mouth.^bConfirmation of young accompanied by adults in the case of waterfowl and rallids.

References

- Allen, D.C., Vaughn, C.C., Kelly, J.F., Cooper, J.T., Engel, M.H., 2012. Bottom-up biodiversity effects increase resource subsidy flux between ecosystems. *Ecology* 93, 2165–2174.
- Atauri, J.A., de Lucio, J.V., 2001. The role of landscape structure in species richness distribution of birds amphibians, reptiles and lepidopterans in Mediterranean landscapes. *Landsc. Ecol.* 16, 147–159.
- Atkinson, P.W., Buckingham, D.L., Morris, A.J., 2004. What factors determine where invertebrate-feeding birds forage in dry agricultural grasslands? In: *Ecology and conservation of lowland farmland birds II: the road to recovery*. Ibis 146 (Suppl. 2), 99–107.
- BTO (The British Trust for Ornithology), 2014. Garden Birdwatch: Our Methods. BTO, Thetford <http://www.bto.org/volunteer-surveys/gbw/taking-part/instructions/how> (02.07.14).
- Baillie, S.R., et al., 2014. BirdTrends 2014: trends in numbers, breeding success and survival for UK breeding birds. Research Report 662. BTO, Thetford. <http://www.bto.org/birdtrends>.
- Baker, D.J., Freeman, S.N., Grice, P.V., Siriwardena, G.M., 2012. Landscape-scale responses of birds to agri-environment management: a test of the English Environmental Stewardship scheme. *J. Appl. Ecol.* 49, 871–882.
- Barker, A.M., 2004. Insects as food for farmland birds: is there a problem? In: van Emden, H.F., Rothschild, M. (Eds.), *Insects and Bird Interactions*. Intercept, Andover, pp. 37–50.
- Bartels, P., Cucherousset, J., Steger, K., Eklöv, P., Tranvik, L.J., Hillebrand, H., 2012. Reciprocal subsidies between freshwater and terrestrial ecosystems structure consumer resource dynamics. *Ecology* 93, 1173–1182.
- Baxter, C., Fausch, V.K.D., Saunders, W.C., 2005. Tangled webs: reciprocal flows of invertebrate prey link streams and riparian zones. *Freshw. Biol.* 50, 201–220.
- Bennett, A.F., Radford, J.Q., Haslem, A., 2006. Properties of land mosaics: implications for nature conservation in agricultural environments. *Biol. Conserv.* 133, 250–264.
- Benton, T.G., Bryant, D.M., Cole, L., Crick, H.P.Q., 2002. Linking agricultural practise to insect and bird populations: a historical study over three decades. *J. Appl. Ecol.* 36, 673–687.
- Benton, T.G., Vickery, J.A., Wilson, J.D., 2003. Farmland biodiversity: is habitat heterogeneity the key? *Trends Ecol. Evol.* 18, 182–188.
- Bibby, C.J., Burgess, N.D., Hill, D.A., 1992. *Bird Census Techniques*. Academic Press, London.
- Biggs, J., Williams, P., Whitfield, M., Nicolet, P., Weatherby, A., 2005. 15 years of pond assessment in Britain: results and lessons learned from the work of Pond Conservation. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 15, 693–714.
- Burel, F., Butet, A., Delettre, Y., Millán de la Peña, N., 2004. Differential response of selected taxa to landscape context and agricultural intensification. *Landsc. Urban Plann.* 67, 195–204.
- Butler, S.J., Vickery, J.A., Norris, K., 2007. Farmland biodiversity and the footprint of agriculture. *Science* 315, 381–384.
- Céréghino, R., Ruggerio, A., Marty, P., Angélibert, S., 2008. Biodiversity and distribution patterns of freshwater invertebrates in farm ponds of a south-western French agricultural landscape. *Hydrobiologia* 597, 43–51.
- Céréghino, R., Boix, D., Cauchie, H.M., Martens, K., Oertli, B., 2014. The ecological role of ponds in a changing world. *Hydrobiologia* 723, 1–6.
- Chamberlain, D.E., Fuller, R.J., Bunce, R.G.H., Duckworth, J.C., Shrubbs, M., 2000. Changes in the abundance of farmland birds in relation to the timing of agricultural intensification in England and Wales. *J. Appl. Ecol.* 37, 771–788.
- Cody, M.L., 1985. *Habitat Selection in Birds*. Academic Press, Orlando.
- Colwell, R.K., 2009. Estimates S: Statistical Estimation of Species Richness and Shared Species from Samples, Version 8.2, User's Guide and Application. <http://purl.oclc.org/estimates>.
- Cresswell, W., 1996. Surprise as a winter hunting strategy in sparrowhawks *Accipiter nisus*, peregrines *Falco peregrinus* and merlins *F. columbarius*. *Ibis* 138, 684–692.
- Crist, T.O., 2003. Partitioning species diversity across landscapes and regions: a hierarchical analysis of α , β and γ diversity. *Am. Nat.* 162, 734–743.
- Davies, B., Biggs, J., Williams, P., Whitfield, M., Nicolet, P., Sear, D., Bray, S., Maund, S., 2008. Comparative biodiversity of aquatic habitats in the European agricultural landscape. *Agric. Ecosyst. Environ.* 125, 1–8.
- Declerck, S., et al., 2006. Ecological characteristics of small farmland ponds: associations with land use practices at multiple spatial scales. *Biol. Conserv.* 131, 523–532.
- Donald, P.F., Evans, A.D., 2006. Habitat connectivity and matrix restoration: the wider implications of agri-environment schemes. *J. Appl. Ecol.* 43, 209–218.
- Donald, P.F., Green, R.E., Heath, M.F., 2001. Agricultural intensification and the collapse of Europe's farmland bird populations. *Proc. R. Soc. B: Biol. Sci.* 268, 25–29.
- Doxa, A., Bas, Y., Paracchini, M.L., Pointereau, P., Terres, J.M., Jiguet, F., 2010. Low intensity agriculture increases farmland bird abundances in France. *J. Appl. Ecol.* 47, 1348–1356.
- Dreyer, J., Townsend, P.A., Hook III, J.C., Hoekman, D., Zanden, M.J.V., Gratton, C., 2015. Quantifying aquatic insect deposition from lake to land. *Ecology* 96, 499–509.
- Eaton, M.A., et al., 2013. The State of the UK's Birds 2013. RSPB, BTO, WWT, NRT, JNCC, NEA and SNH, Sandy, Bedfordshire.
- Fahrig, L., Brotons, L., Burel, F.G., Crist, T.O., Fuller, R.J., Sirami, C., Siriwardena, G.M., Martin, J.L., 2011. Functional landscape heterogeneity and animal biodiversity in agricultural landscapes. *Ecol. Lett.* 14, 101–112.
- Florencio, M., Diaz-Paniagua, C., Gomez-Rodriguez, C., Serrano, L., 2014. Biodiversity patterns in a macroinvertebrate community of a temporary pond network. *Insect Conserv. Diver.* 7, 4–21.
- Foley, J.A., DeFries, R., Asner, G.P., 2005. Global consequences of land use. *Science* 309, 570–574.
- Ford, H.A., Barrett, G.W., Saunders, D.A., Recher, H.F., 2001. Why have birds in the woodlands of Southern Australia declined? *Biol. Conserv.* 97, 71–88.
- Froneman, A., Mangnall, J., Little, R.M., Crowe, T.M., 2001. Water-bird assemblages and associated habitat characteristics of farm ponds in the Western Cape, South Africa. *Biodivers. Conserv.* 10, 251–270.
- Fuller, R.J., Gregory, R.D., Gibbons, D.W., Marchant, J.H., Wilson, J.D., Baillie, S.R., Carter, N., 1995. Population declines and range contractions among lowland farmland birds in Britain. *Conserv. Biol.* 9, 1425–1441.
- Fuller, R.J., 2000. Relationships between recent changes in lowland British agriculture and farmland bird populations: an overview. In: Aebischer, N.J., Evans, A.D., Grice, P., Vickery, J.A. (Eds.), *The Conservation and Ecology of Lowland Farmland Birds*. Tring, British Ornithologist's Union, pp. 5–16.
- Gee, J.H.R., Smith, B.D., Lee, K.M., Griffiths, S.W., 1997. The ecological basis of freshwater pond management for biodiversity. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 7, 91–104.
- Gregory, R.D., Noble, D.G., Custance, J., 2004. The state of play of farmland birds: population trends and conservation status of lowland farmland birds in the United Kingdom. *Ibis* 146, 1–13.
- Hinden, H., Oertli, B., Menetrey, N., Sager, L., Lachavanne, J.B., 2005. Alpine pond biodiversity: what are the related environmental variables? *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 15, 613–624.
- Hinsley, S.A., Bellamy, P.E., 2000. The influence of hedge structure, management and landscape context on the value of hedgerows to birds: a review. *J. Environ. Manage.* 60, 33–49.
- Holland, J.M., 2004. The impact of agriculture and some solutions for arthropods and birds. In: van Emden, H.F., Rothschild, M. (Eds.), *Insects and Bird Interactions*. Intercept, Andover, pp. 51–73.
- Hurlbert, S.H., 1971. The nonconcept of species diversity: a critique and alternative parameters. *Ecology* 52, 577–586.
- IBM Corp., 2011. IBM SPSS Statistics for Windows Version 20.0. IBM Corp, Armonk, New York.
- Joint Nature Conservation Committee (JNCC), 2007. Report on the Species and Habitat Review. Report by the Biodiversity Reporting and Information Group (BRIG) to the UK Standing Committee June 2007.
- Josefsson, J., Berg, A., Hiron, M., Pärta, T., Eggers, S., 2013. Grass buffer strips benefit invertebrate and breeding skylark numbers in a heterogeneous agricultural landscape. *Agric. Ecosyst. Environ.* 181, 101–107.
- Kleijn, D., et al., 2006. Mixed biodiversity benefits of agri-environment schemes in five European countries. *Ecol. Lett.* 9, 243–254.
- Kleijn, D., Rundlöf, M., Scheper, J., Smith, H.G., Tscharntke, T., 2011. Does conservation on farmland contribute to halting the biodiversity declines? *Trends Ecol. Evol.* 26, 474–481.
- Knight, T.M., McCoy, M.W., Chase, J.M., McCoy, K.A., Holt, R.D., 2005. Trophic cascades across ecosystems. *Nature* 437, 880–883.
- Lawton, J.H., et al., 2010. Making Space For Nature: a review of England's wildlife sites and ecological network. Report to Defra.
- Le Viol, I., Jiguet, F., Brotons, L., Herrando, S., Lindström, Å., Pearce-Higgins, J.W., Reif, J., Van Turnhout, C., Devictor, V., 2012. More and more generalists: two decades of changes in the European avifauna. *Biol. Lett.* 8, 780–782.
- Lemmens, P., Mergeay, J., De Bie, T., Van Wichelen, J., De Meester, L., Declerck, S.A.J., 2013. How to maximally support local and regional biodiversity in applied conservation? Insights from pond management. *PLoS One* 8, e72538.
- Marja, R., Herzog, I., 2012. The importance of drainage ditches for farmland birds in agricultural landscapes in the Baltic countries: does field type matter? *Ornis Fenn.* 89, 1–12.
- Matuszak, A., Moertl, M., Quillfeldt, P., Bauer, H.G., 2014. Macrophyte-associated macroinvertebrates as an important food source for wintering waterbirds at Lake Constance. *Limnology* 15, 69–76.
- McAbendroth, L., Ramsay, P.M., Foggo, T., Rundle, S.D., Bilton, D.T., Persson, L., 2005. Does macrophyte fractal complexity drive invertebrate diversity, biomass and body size distributions? *Oikos* 111, 279–290.
- McCracken, D.L., Tallwin, J.R., 2004. Swards and structure: the interactions between farming practises and bird food resources in lowland grasslands. *Ibis* 146, 170–181.
- McKinstry, M.C., Anderson, S.H., 2002. Creating wetlands for waterfowl in Wyoming. *Ecol. Eng.* 18, 293–304.
- Natural England, 2015. Countryside Stewardship Handbook – November 2015.
- Neuschultz, E.L., Brown, M., Farwig, N., 2013. Frequent bird movements across a highly fragmented landscape: the role of species traits and forest matrix. *Anim. Conserv.* 16, 170–179.
- Newton, I., 1998. *Population Limitation in Birds*. Academic Press, San Diego.
- Palmer, M.A., Bell, S.L., Butterfield, I.A., 1992. A botanical classification of standing waters in Britain: an application for conservation and monitoring. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 2, 125–143.
- Perkins, A.J., Whittingham, M.J., Bradbury, R.B., Wilson, J.D., Morris, A.J., Barnett, P.R., 2000. Habitat characteristics affecting use of lowland agricultural grassland by birds in winter. *Biol. Conserv.* 95, 279–294.
- Pino, J., Rodà, F.J., Ribas, J., Pons, X., 2000. Landscape structure and bird species richness: implications for conservation in rural areas between natural parks. *Landsc. Urban Plann.* 49, 35–48.

- Pisces Conservation Ltd, 2002. Species Diversity and Richness. Version 3.02. Pisces Conservation Ltd, Lymington.
- Prince, H.C., 1964. The origin of pits and depressions in Norfolk. *Geography* 49, 15–32.
- R Core Team, 2015. R: A Language and Environment for Statistical Computing, Version 3.2.2. R Foundation for Statistical Computing, Vienna, Austria.
- Richardson, J.S., Zhang, Y., Marczak, L.B., 2010. Resource subsidies across the land-freshwater interface and responses in recipient communities. *River Res. Appl.* 26, 55–66.
- Rioux Paquette, S., Pelletier, F., Garant, D., Bélisle, M., 2014. Severe recent decrease of adult body mass in a declining insectivorous bird population. *Proc. R. Soc. B: Biol. Sci.* 282, 20140649.
- Robinson, R.A., Sutherland, W.J., 2002. Post-war changes in arable farming and biodiversity in Great Britain. *J. Appl. Ecol.* 39, 157–176.
- Robinson, R.A., Hart, J.D., Holland, J.M., Parrott, D., 2004. Habitat use by seed-eating birds: a scale-dependent approach. *Ibis* 146, 87–98.
- Ruggiero, A., Céréghino, R., Figuerola, J., Marty, P., Angélibert, S., 2008. Farm ponds make a contribution to the biodiversity of aquatic insects in a French agricultural landscape. *C. R. Biol.* 331, 298–308.
- Sabatier, R., Doyen, L., Tichit, M., 2014. Heterogeneity and the trade-off between ecological and productive functions of agro-landscapes: a model of cattle-bird interactions in a grassland agroecosystem. *Agric. Syst.* 126, 38–49.
- Santoul, F., Gaujard, A., Angélibert, S., Mastroiello, S., Céréghino, R., 2009. Gravel pits support water-bird diversity in an urban landscape. *Hydrobiologia* 634, 107–114.
- Sayer, C.D., Copp, G.H., Emson, D., Godard, M.J., Zieba, G., Wesley, K.J., 2011. Towards the conservation of crucian carp *Carassius carassius*: understanding the extent and causes of decline within part of its native English range. *J. Fish Biol.* 79, 1608–1624.
- Sayer, C.D., Andrews, K., Shilland, E., Edmonds, N., Edmonds-Brown, R., Patmore, I.R., Emson, D., Axmacher, J.C., 2012. The role of pond management for biodiversity conservation in an agricultural landscape. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 22, 626–638.
- Sayer, C.D., Shilland, E., Greaves, H., Dawson, B., Patmore, I.R., Emson, D., Alderton, E., Robinson, P., Andrews, K., Axmacher, J.A., Wiik, E., 2013. Managing British ponds – conservation lessons from a Norfolk farm. *Br. Wildl.* 25 (1), 21–28.
- Scherr, S.J., McNeely, J.A., 2008. Biodiversity conservation and agricultural sustainability: towards a new paradigm of 'ecoagriculture' landscapes. *Phil. Trans. R. Soc. B: Biol. Sci.* 363, 477–494.
- Schummer, M.L., Palframan, J., McNaughton, E., Barney, T., Petrie, S.A., 2012. Comparisons of bird, aquatic macroinvertebrate and plant communities among dredged ponds and natural wetland habitats at Long Point Lake Erie, Ontario. *Wetlands* 32, 945–953.
- Sebastián-González, E., Sánchez-Zapata, J.A., Botella, F., 2010. Agricultural ponds as alternative habitat for waterbirds: spatial and temporal patterns of abundance and management strategies. *Eur. J. Wildl. Res.* 56, 11–20.
- Siriwardena, G.M., Calbrade, N.A., Vickery, J.A., 2008. Farmland birds and late winter food: does seed supply fail to meet demand? *Ibis* 150, 585–595.
- Soininen, J., Bartels, P., Heino, J., Luoto, M., Hillebrand, H., 2015. Toward more integrated ecosystem research in aquatic and terrestrial environments. *Bioscience* 65, 174–182.
- Sparks, T.H., Parish, T., Hinsley, S.A., 1996. Breeding birds in field boundaries in an agricultural landscape. *Agric. Ecosyst. Environ.* 60, 1–8.
- Stenroth, K., Polvi, L.E., Fältström, E., Jonsson, M., 2015. Land-use effects on terrestrial consumers through changed size structure of aquatic insects. *Freshw. Biol.* 60, 136–149.
- Stoate, C., Báldi, A., Beja, P., Boatman, N.D., Herzin, I., van Doorn, A., de Snoo, G.R., Rakosy, L., Ramwell, C., 2009. Ecological impacts of early 21st century agricultural change in Europe – a review. *J. Environ. Manage.* 91, 22–46.
- ter Braak, C., Smilauer, P., 2002. CANOCO Reference Manual and CanoDraw for Windows User's Guide: Software for Canonical Community Ordination (version 4.5). Microcomputer Power, Ithaca, New York.
- Thomaz, S.M., da Cunha, E.R., 2010. The role of macrophytes in habitat structuring in aquatic ecosystems: methods of measurement, causes and consequences on animal assemblages' composition and biodiversity. *Acta Limnol. Bras.* 22, 218–236.
- Toms, M., 2004. Garden Birdwatch Handbook. British Trust for Ornithology, BTO, Thetford.
- Tscharntke, T., Klein, A.M., Kruess, A., Steffan-Dewenter, I., Thies, C., 2005. Landscape perspectives on agricultural intensification and biodiversity: ecosystem service management. *Ecol. Lett.* 8, 857–874.
- van Zanten, B.T., et al., 2014. European agricultural landscapes: common agricultural policy and ecosystem services: a review. *Agron. Sustain. Dev.* 34, 309–325.
- Weibull, A.C., Östman, O., Granqvist, A., 2003. Species richness in agroecosystems: the effect of landscape: habitat and farm management. *Biodivers. Conserv.* 12, 1335–1355.
- Welsh Government, 2015. Glastir Entry Booklet 2: Technical Guidance 2015. .
- Whittingham, M.J., Krebs, J.R., Swetnam, R.D., Thewlis, R.M., Wilson, J.D., Freckleton, R.P., 2009. Habitat associations of British breeding farmland birds. *Bird Study* 56, 43–52.
- Williams, P., Whitfield, M., Biggs, J., Bray, S., Fox, G., Nicolet, P., Sear, D., 2004. Comparative biodiversity of rivers streams, ditches and ponds in an agricultural landscape in Southern England. *Biol. Conserv.* 115, 329–341.
- Williams, P., Biggs, J., Nicolet, P., 2010. New clean-water ponds – a way to protect freshwater biodiversity. *Br. Wildl.* 22, 77–85.
- Wood, P.J., Greenwood, M.T., Agnew, M.D., 2003. Pond biodiversity and habitat loss in the UK. *Area* 35, 206–216.